

1/9




FIG.1

2/9




FIG.2

3/9

EFFECT OF 2'-O-METHYL GAPPED OLIGOS





FIG.3

4/9

**FIG. 4**

5/9





FIG. 5

6/9

FIG. 6

PLASMA CONCENTRATION OF ISIS 3082, AND 2' O-PROPYL ANALOGS
(ISIS 9045 AND 9046) IN BALB/c MICE AFTER INTRAVENOUS INJECTION



7/9

**ISIS 3082 TISSUE DISTRIBUTION IN BALB/c MICE
AS PERCENT OF DOSE AT 1, 3, 8 AND 24 HOURS AFTER
INTRAVENOUS INJECTION**




FIG. 7

8/9

**ISIS 9045 (2' O-PROPYL PHOSPHOROTHIAE ANALOG OF ISIS 3082)
TISSUE DISTRIBUTION IN BALB/c MICE
AS PERCENT OF DOSE AT 1, 3, 8 AND 24 HOURS AFTER
INTRAVENOUS INJECTION**




FIG.8

9/9

**TISSUE DISTRIBUTION OF ISIS 9046 (2' O-PROPYL GAPMER) IN BALB/c MICE
AT 1, 3, 8 AND 24 HOURS AFTER INTRAVENOUS INJECTION**




FIG.9